

Excited state spectra at the superfluid-insulator transition out of paired condensates

Stephen Powell

Department of Physics, Yale University, New Haven, CT 06520-8120

Subir Sachdev

Department of Physics, Harvard University, Cambridge MA 02138

(Dated: August 28, 2006)

We describe gapped single-particle and collective excitations across a superfluid to insulator quantum phase transition of particles (bosons or fermions) in a periodic potential, with an even number of particles per unit cell. We demonstrate that the dynamics is controlled by a *quantum impurity* problem of a localized particle interacting with the bulk critical modes. Critical exponents are determined by a renormalization group analysis. We discuss applications to spin oscillations of ultracold atoms in optical lattices, and to the electronic phases in the cuprate and related compounds.

PACS numbers: 05.30.-d, 03.75.Kk, 71.10.-w

A number of experiments have observed coherent spin oscillations of trapped, ultracold spinor bosonic atoms. With a superfluid ground state [1, 2, 3, 4, 5, 6], these oscillations are well described by the classical (Gross-Pitaevski) equations which control the time evolution of the multi-component atomic condensate. Recent experiments [7, 8] have also observed oscillations in a Mott insulating state obtained by placing the atoms in an optical lattice; in this situation, the spin oscillations can be viewed as the Rabi precession [7, 9] between sharp quantum states localized within each minimum of the optical lattice.

In this paper, we investigate the connection between these two disparate pictures of the spin oscillations, by describing ground state spin correlations at the quantum critical point separating the superfluid and insulating phases. Bosonic atoms with total spin $F = 1$ and $F = 2$ display a remarkably rich variety of superfluid and insulating phases [10, 11, 12, 13, 14], and have a corresponding plethora of universality classes of quantum phase transitions separating such phases. Superfluid-insulator transitions are also possible for paired fermionic atoms in an optical lattice. We defer a more complete classification of the spin dynamics at such transitions to a forthcoming paper. Here we focus on a class, with an even integer number of particles per unit cell, which displays non-trivial collective behavior induced by a strong coupling between the spin excitations and the critical number and phase fluctuations of the superfluid-insulator transition; the class includes both the fermionic and bosonic cases. The problem is mapped exactly onto a quantum ‘impurity’ problem, which couples a single localized spin excitation (the ‘impurity’) to the bulk critical modes; a solution using the renormalization group yields new critical exponents and scaling functions. Our theory has some analogies to simpler models of the Kondo and X-ray edge effect in metals, and opens the way towards observing strongly-coupled quantum impurity physics in

ultracold atom systems.

Our results also apply to the superfluid-insulator transition in electronic systems, with an even integer number of electrons per unit cell. Such a situation can arise in the cuprate or related compounds, with a periodic potential generated spontaneously by ‘stripe/checkerboard’ or charge density wave order: recent experiments in insulating spin ladder compounds [15] have shown that each unit cell contains a pair of holes. Our results predict the frequency dependence of the electron photoemission spectrum across a superfluid-insulator transition in which the ‘stripe’ order is present on both sides of the transition; in other words, for a transition between a modulated insulator and a supersolid. The predictions of the spectrum are for gapped single particle excitations at positions of the gap maxima or minima in the Brillouin zone *e.g.* at the analog of the ‘antinodal’ points.

We will study superfluid-insulator transitions at which the energy gap to both single-particle and spin excitations remains nonzero at the transition. The superfluid order parameter for the transition is then necessarily a spin singlet. The order parameter also carries a nonzero particle number, or ‘charge’ Q . Here, and henceforth, ‘particle’ refers to either a single ultracold bosonic or fermionic atom, or an electron (but *not* a Cooper pair). For $F \neq 0$ particles, this means that the simplest case has an order parameter, Ψ , with $Q = 2$ and $F = 0$, corresponding to the annihilation operator for a spin-singlet pair of particles, *e.g.* a Cooper pair.

For definiteness, we will develop our results in the context of a Bose-Hubbard model for $F \neq 0$ bosons in an optical lattice, and indicate the generalization to the fermionic case later. For this model, we consider the transition from a ‘spin-singlet insulator’ (SSI), a Mott insulator with an even number of atoms per lattice site and no spin order, to a ‘spin-singlet condensate’ (SSC), in which singlet pairs of bosons have condensed, but there is no single boson condensate. We will begin with a simple

mean-field theory of the Bose-Hubbard model, and then turn to a field theory of the critical properties of this and the corresponding fermionic model.

The lattice bosons are annihilated with operators $a_{i,m}$ on lattice site i and spin projection $m = -F \dots F$. The Bose-Hubbard Hamiltonian can then be written as $\mathcal{H} = -tT + V$, where T is the kinetic energy term,

$$T = \sum_{\langle i,j \rangle, m} (a_{i,m}^\dagger a_{j,m} + \text{h.c.}), \quad (1)$$

and V is the on-site interaction:

$$V = \sum_i [U(\hat{n}_i - N)^2 + J|\mathbf{L}_i|^2], \quad (2)$$

where $\hat{n}_i = \sum_m a_m^\dagger a_m$ is the boson number operator on site i , and \mathbf{L}_i is the total spin operator on site i . We have made the spin-independent part of the interaction explicitly symmetric around N particles per site. For $F = 1$, the final term is the most general spin-dependent interaction, but further terms are necessary for higher spin. To favor spin-singlet pairing in the ground state, we require N to be even, and $J > 0$.

In the case when $t = 0$, the Hamiltonian is simply a sum of terms acting on a single site, containing only the commuting operators \hat{n} and $|\mathbf{L}|^2$. The ground state on each site is therefore a spin singlet of N bosons.

An appropriate mean-field Hamiltonian is $\mathcal{H}_{\text{mf}} = V - T_\psi - T_\Psi - T_\Phi$, where V is the same on-site interaction as in Eq. (2). T_ψ is the standard mean-field decoupling of the hopping term, generalized to the case with spin,

$$T_\psi = \sum_i [\psi_m a_{i,m}^\dagger + \psi_m^* a_{i,m}], \quad (3)$$

where ψ_m is a (c-number) constant vector, which will be used as a variational parameter. The remaining terms allow for the possibility of a spin-singlet condensate through the parameters Ψ and Φ :

$$T_\Psi = \sum_{i,m} (-1)^{F+m} [\Psi a_{i,m}^\dagger a_{i,-m}^\dagger + \Psi^* a_{i,m} a_{i,-m}], \quad (4)$$

and

$$T_\Phi = \sum_{\langle i,j \rangle, m} (-1)^{F+m} [\Phi a_{i,m}^\dagger a_{j,-m}^\dagger + \Phi^* a_{i,m} a_{j,-m}], \quad (5)$$

where the factors of $(-1)^{F+m}$ are Clebsch-Gordan coefficients that cause the boson operators to form spin-singlet pairs.

We now use the ground state of \mathcal{H}_{mf} , which we denote $|\text{mf}\rangle$, as a variational ansatz and define

$$E_{\text{mf}}(\psi_m, \Psi, \Phi) = \langle \text{mf} | \mathcal{H} | \text{mf} \rangle, \quad (6)$$

which should be minimized by varying the three parameters. If this minimum occurs for vanishing values of all




FIG. 1: Phase diagram calculated using mean-field theory. The calculation has been performed for spin $F = 1$ and filling factor $N = 2$. The horizontal and vertical axes give the tunneling strength t and the spin-dependent interaction J , both in units of the spin-independent part of the interaction, U .

three parameters, then $|\text{mf}\rangle$ breaks no symmetries and the Mott insulator is favored. A nonzero value for ψ_m at the minimum corresponds to a simple ‘polar condensate’ (PC), breaking spin-rotation symmetry; vanishing ψ_m but nonzero values of Ψ and/or Φ corresponds to a paired spin-singlet condensate.

Since \mathcal{H}_{mf} contains terms (within T_Φ) that link adjacent sites, it cannot be straightforwardly diagonalized, as in the standard mean-field theory for the spinless Bose-Hubbard model. To find the phase boundaries, however, we need only terms up to quadratic order in the variational parameters, which can be found using perturbation theory. Figure 1 shows the phase boundaries so obtained.

We now turn to an analysis of the correlation functions of gapped spin-carrying modes across the SSI/SSC transition. We will assume we are in a regime in which the lowest excitation with a nonzero spin is created by an elementary boson operator, ψ_m , with $Q = 1$ and $F \neq 0$, with $m = -F, \dots, F$.

General symmetry arguments can be used to deduce the low-energy theory coupling Ψ to ψ_m in the vicinity of the superfluid-insulator transition. For the Ψ field alone, we have the familiar $|\Psi|^4$ field theory of the superfluid-insulator transition,

$$\mathcal{S}_\Psi = \int d^d \mathbf{x} d\tau \left(|\partial \Psi|^2 + r_\Psi |\Psi|^2 + \frac{u_\Psi}{2} |\Psi|^4 + \dots \right) \quad (7)$$

in d spatial dimensions with co-ordinate \mathbf{x} , imaginary time τ , and $(d+1)$ -dimensional derivative $\partial^2 = \partial_\tau^2 + \nabla^2$. The even integer number of bosons per lattice site ensures that there is particle-hole symmetry in \mathcal{S}_Ψ in the vicinity of the critical point.

The same particle-hole symmetry applies to the gapped ψ_m field, but it is convenient to treat separately the particle-like (with $Q = 1$) and hole-like (with $Q = -1$)

Since ψ_m has only gapped excitations, it has no effect on the bulk scaling, which is therefore given by the standard results for a complex $|\Psi|^4$ theory. Rescaling by the standard phase space factor $\hat{u} = u_\Psi/S_{d+1}$ (where $S_d = 2/(\Gamma(d/2)(4\pi)^{d/2})$), there is a fixed point with coupling $\hat{u} = \epsilon/5 + \mathcal{O}(\epsilon^2)$ in $3 - \epsilon$ dimensions, at which the scaling dimension of the field is given by $[\Psi] = 1 - \frac{\epsilon}{2} + \epsilon^2/100 + \mathcal{O}(\epsilon^3)$.

These results for the bulk, along with the self-energy diagram above, when extended to two loops, lead to the RG flow equation for the coupling to the ‘impurity’:

$$\frac{d\hat{g}}{d\ell} = \frac{\epsilon}{2}\hat{g} - \hat{g}^3 + 2\hat{g}^5 - \frac{\hat{u}^2\hat{g}}{4} - \frac{2\pi^2\hat{u}\hat{g}^3}{3} + \mathcal{O}(\hat{g}(\hat{u}, \hat{g}^2)^3), \quad (11)$$

where we have defined $\hat{g} = g(4\pi)^{(d+1)/2}/\Gamma((d-1)/2)$. The coupling therefore approaches a fixed-point value with $\hat{g}^2 = \epsilon/2 - (\pi^2/15 - 49/100)\epsilon^2 + \mathcal{O}(\epsilon^3)$, so that the perturbative expansion at this point is indeed equivalent to an expansion in ϵ .

Finally, using the wavefunction renormalization of ψ_m , we arrive at the rescaling of the Green function G^ψ , which obeys

$$G^\psi(\lambda + \delta\omega) = e^{-y\ell} G^\psi(\lambda + e^{-\ell}\delta\omega), \quad (12)$$

where $y = 1 - \hat{g}^2 + \hat{g}^4 + \mathcal{O}(\hat{g}^6)$, with \hat{g} set to the fixed point of Eq. (11); this yields $y = 1 - \epsilon/2 + (6/25 - \pi^2/15)\epsilon^2 + \mathcal{O}(\epsilon^3)$. This can be iterated to give $G^\psi(\lambda + \delta\omega) \sim \delta\omega^{-y}$. Note that, at least to this order, $y < 1$, so that the quasi-particle pole at $\omega = \lambda$ is replaced by a power-law threshold singularity. This exponent, determining the spectral density of the ‘photoemission’ of a single hole or particle at a band minimum or maximum, is one of our central results.

A similar calculation applies to the two-particle threshold singularity at $\omega = 2\lambda$. This is associated with the renormalization of the $T_{mm'}p_m h_{m'}$ correlation function, where $T_{mm'}$ is an arbitrary matrix. It is then also necessary to calculate the following insertion diagram (at one loop order):

$$H_1 = \dots \quad (13)$$

which accounts for the renormalization associated with bringing the particle and hole operators to the same space-time point. The resulting calculation is very similar and leads to the result for the two-particle propagator

$$\Pi(\omega, \mathbf{k} = 0) \sim (2\lambda - \omega)^{-y_2}, \quad (14)$$

where the exponent y_2 depends upon whether the matrix $T_{mm'}$ is symmetric or antisymmetric. For bosons, a symmetric T has $y_2 = 1 - 2\hat{g}^2 + \mathcal{O}(\hat{g}^6)$, while an antisymmetric T has $y_2 = 1$ exactly; these exponents therefore determine the threshold singularities for excitations with even and odd total spin F , respectively. For fermions, the same results apply, but with the roles of symmetric/antisymmetric T reversed.

This paper has presented results for a variety of threshold singularities in the spectral functions of a system undergoing a superfluid-insulator transition, with an even number of particles per unit cell. This has direct application to experiments on ultracold atoms and to the cuprate compounds. It would be interesting to extend these methods to study non-equilibrium phenomena, such as those measured in Ref. [7], by methods explored in recent studies of non-equilibrium quantum criticality [18, 19] (which have so far been limited to 1+1 dimensions).

We thank E. Demler and R. Shankar for useful discussions. This research was supported by the NSF grants DMR-0537077 and DMR-0342157.

-
- [1] J. Stenger *et al.*, Nature **396**, 345 (1998).
 - [2] M. S. Chang *et al.*, Phys. Rev. Lett. **92**, 140403 (2004).
 - [3] J. M. Higbie *et al.*, Phys. Rev. Lett. **95**, 050401 (2005).
 - [4] M. S. Chang *et al.*, Nature Phys. **1**, 111 (2005).
 - [5] J. Mur-Petit *et al.*, Phys. Rev. A **73**, 013629 (2006).
 - [6] J. Kronjäger *et al.*, cond-mat/0509083.
 - [7] A. Widera *et al.*, Phys. Rev. Lett. **95**, 190405 (2005).
 - [8] F. Gerbier *et al.*, Phys. Rev. A **73**, 041602 (2006).
 - [9] H.-J. Huang and G.-M. Zhang, cond-mat/0601188.
 - [10] A. Imambekov, M. Lukin, and E. Demler, Phys. Rev. A **68**, 063602 (2003).
 - [11] S. Tsuchiya, S. Kurihara, and T. Kimura, Phys. Rev. A **70**, 043628 (2004).
 - [12] D. Rossini *et al.*, J. Phys. B: At. Mol. Opt. Phys. **39**, S163 (2006).
 - [13] L. Zawitkowski *et al.*, cond-mat/0603273.
 - [14] R. Barnett, A. Turner, and E. Demler, cond-mat/0607253.
 - [15] A. Rusydi *et al.*, Phys. Rev. Lett. **97**, 016403 (2006).
 - [16] S. Sachdev, M. Troyer, and M. Vojta, Phys. Rev. Lett. **86**, 2617 (2001).
 - [17] S. Sachdev, Physica C **357**, 78 (2001).
 - [18] K. Sengupta, S. Powell, and S. Sachdev, Phys. Rev. A **69**, 053616 (2004).
 - [19] P. Calabrese and J. Cardy, Phys. Rev. Lett. **96**, 136801 (2006).