

Spin-Peierls ground states of the quantum dimer model: A finite-size study

Subir Sachdev

Center for Theoretical Physics, P.O. Box 6666, Yale University, New Haven, Connecticut 06511

Institute for Theoretical Physics, University of California, Santa Barbara, California 93106

(Received 10 March 1989)

Exact diagonalization results on the quantum dimer Hamiltonian of Rokhsar and Kivelson on lattice sizes up to 6×6 are presented. Correlation functions of a new order parameter characterizing the crystalline states of the model are calculated. All of the results are consistent with the existence of a direct transition between a columnar crystalline state and a staggered crystalline state with no intermediate spin-fluid state. However, the existence of a spin-fluid state over a small, but finite, range of parameters has not been definitively ruled out.

INTRODUCTION

The quantum dimer model was introduced recently by Rokhsar and Kivelson¹ (RK) as a phenomenological description of a non-Néel phase of the spin- $\frac{1}{2}$ Heisenberg quantum antiferromagnet on a square lattice. Interest in the latter model is motivated partly by the suggestion of Anderson² that properties of the non-Néel phase are important in understanding the occurrence of high temperature superconductivity in $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$ and $\text{YBa}_2\text{Cu}_3\text{O}_{7-x}$.³ In a phase with exponentially decaying two-spin correlation functions, RK argued that it was a reasonable approximation to truncate the Hilbert space to states that can be expressed as a tensor product of nearest-neighbor singlet pairs of spins. There is clearly a one-to-one mapping between such states and the set of all close-packed dimer coverings of the square lattice. RK also argued that the effects of the nonorthogonality of these states could be absorbed into a redefinition of the parameters in the following phenomenological Hamiltonian H acting upon an orthogonal basis set of states represented by all the different close-packed dimer coverings of the square lattice:

$$H = -J \sum_{\square} |1\rangle\langle 2| + \text{H.c.} + V \sum_{\square} |1\rangle\langle 1| + V \sum_{\square} |2\rangle\langle 2| \quad (1)$$

The dark lines denote dimers on the links of the square lattice and all the sums extend over all the elementary plaquettes of the square lattice. The first term is a dimer kinetic energy and the last two terms are the diagonal potential energies. Read and Sachdev⁴ have recently shown that H may be obtained in a formal $1/N$ expansion of a particular nearest-neighbor $\text{SU}(N)$ antiferromagnet with the states at each site transforming with an appropriately chosen representation of $\text{SU}(N)$, and exchange constant J_s ; at order $1/N$ the parameters obtained in this expansion are $J = 2J_s/N$ and $V = 0$. We also have the correspondence $n_d(\mathbf{r} + \hat{\epsilon}/2) \sim -\mathbf{S}(\mathbf{r}) \cdot \mathbf{S}(\mathbf{r} + \hat{\epsilon})$ between the dimer number operators $n_d(\mathbf{R})$ and the spin- $\frac{1}{2}$ generators \mathbf{S} of the Heisenberg model ($\hat{\epsilon}$ is an elementary link of the square lattice). The symbol \sim implies that correlation functions of n_d in the ground state of the dimer Hamiltonian H should have the same asymptotic form as the

corresponding correlation functions of $\mathbf{S}(\mathbf{r}) \cdot \mathbf{S}(\mathbf{r} + \hat{\epsilon})$ in the non-Néel phase of the Heisenberg antiferromagnet.

In a separate development, Read and Sachdev^{4,5} have shown, using several different semiclassical and $1/N$ expansions of $\text{SU}(N)$ antiferromagnets, that the non-Néel phase exhibits spin-Peierls or valence-bond solid order. For the quantum-dimer model, these phases correspond to a crystallization of the dimer bosons into a "column" state with the symmetry of Fig. 1(a); note that this state is fourfold degenerate and completely breaks the Z_4 lattice rotational symmetry.

We briefly review known results and open questions on the properties of H :

(i) The dimer states are characterized by conserved winding numbers (N_x, N_y) (Ref. 1) (specified below) with $-L/2 < N_x, N_y < L/2$ on a $L \times L$ lattice.

(ii) For $V > J$ the exact ground state is the fourfold degenerate "staggered" crystalline state¹ shown in Fig. 1(b). These states have winding numbers $(\pm L/2, 0)$ and $(0, \pm L/2)$.

(iii) At exactly $V = J$, the lowest energy states in all the




FIG. 1. (a) The perfectly ordered column state and (b) the staggered state; this state is the ground state of H for $V > J$. (c) The phase factors θ and (d) ζ on the links of the lattice.

winding-number sectors are degenerate; these states are equal superpositions of all dimer coverings with a fixed winding number.¹ Correlation functions in the equal-superposition state on a square lattice with free boundary conditions [this restricts the state to be in the (0,0) winding-number sector] can be exactly calculated using the results of Fisher and Stephenson,⁶ we shall refer to this particular eigenstate as the Fisher-Stephenson (FS) state. The FS state has no broken lattice symmetry: correlations of the column state order parameter, introduced below, fall off as $1/r^2$.⁷

(iv) For $V < \kappa J$, where $\kappa \leq 1$ is an unknown constant, the ground state of H breaks lattice rotational symmetry and crystallizes into a state with the symmetry of the column state shown in Fig. 1(a). Thus, there is a possible range of values $\kappa J \leq V \leq J$ over which H displays a spin-fluid ground state. In fact, RK suggested that $\kappa < 1$ and that the spin fluid was thus stable over a finite range of V .

The primary purpose of this paper is to obtain more information on the value of κ by exact diagonalization of H on a $L \times L$ lattice with periodic boundary conditions with sizes up to $L=6$. Our results allow us to conclude with reasonable certainty that $\kappa > 0.5$, although all of our results are consistent with a value $\kappa=1$. Thus for the parameters obtained in the $1/N$ expansion ($V=0$), we find a crystalline ground state.

It is useful at this juncture to introduce the complex order parameters $\Psi_{\text{col}}^p(\mathbf{r})$ to measure the breaking of rotational symmetry in the column phase

$$\Psi_{\text{col}}^p(\mathbf{r}) = \sum_{\hat{\mathbf{e}}} [\theta(\mathbf{r} + \hat{\mathbf{e}}/2)]^p n_d(\mathbf{r} + \hat{\mathbf{e}}/2), \quad (2)$$

where, as before, the sum over $\hat{\mathbf{e}}$ extends over $\hat{\mathbf{x}}, -\hat{\mathbf{x}}, \hat{\mathbf{y}},$ and $-\hat{\mathbf{y}}$; the $\theta(\mathbf{R})$ take the fixed values 1, i , -1 , and $-i$ on the links of the square lattice, as shown in Fig. 1(c), and are chosen such that under a rotation by $n\pi/2$ about any point on the A sublattice $\Psi_{\text{col}}^p \rightarrow e^{in\pi p/2} \Psi_{\text{col}}^p$, while on the B sublattice $\Psi_{\text{col}}^p \rightarrow e^{-in\pi p/2} \Psi_{\text{col}}^p$. These properties imply that $\langle \Psi_{\text{col}}^p \rangle = 0$ for $p \neq (\text{mod} 4)$ in any spin-fluid state with unbroken lattice rotational symmetry while $|\langle \Psi_{\text{col}}^p \rangle| \neq 0$ for $p \neq (\text{mod} 4)$ in the column phase.

The order parameter associated with the staggered state is

$$\Psi_{\text{st}}^p(\mathbf{r}) = \sum_{\hat{\mathbf{e}}} [\zeta(\mathbf{r} + \hat{\mathbf{e}}/2)]^p n_d(\mathbf{r} + \hat{\mathbf{e}}/2), \quad (3)$$

where $\zeta(\mathbf{R})$ takes the values shown in Fig. 1(d). Note that $\Psi_{\text{col}}^2 = \Psi_{\text{st}}^2$. It can also be shown that $\sum_{\mathbf{r} \in A} \Psi_{\text{st}}^1(\mathbf{r}) = L(N_y + iN_x)$, where the summation over \mathbf{r} extends over the A sublattice, and N_x and N_y are the conserved winding numbers.¹ The connection of Ψ_{st} with the conserved winding numbers will restrict its utility in the finite-size calculations presented below.

NUMERICAL RESULTS

With momentum and winding-number conservation the calculation required diagonalization of matrices of dimensions up to 1256: the diagonalization was carried out by a modified Lanczos method.⁸ We note that an analogous procedure on a 8×8 lattice would require diagonalization

of matrices of order 10^6 , which is clearly prohibitive. For all $V < J$ the ground state of H was found in the winding number $(N_x, N_y) = (0, 0)$ sector. All of the correlation functions presented below are obtained with the Hilbert space restricted to the (0,0) winding-number sector for all values of V . For $V < J$, this implies that, for large enough system sizes, the low-energy states will consist of domains of the staggered states in different orientations; the domains are chosen in a manner that restricts the global winding number to (0,0). The order parameter correlation functions are trivially calculable in the exact staggered ground state [Fig. 1(b)] for $V > J$ and do not yield any useful information.

We begin by measuring the correlation function

$$\chi^1 = \frac{1}{L^2} \left\langle \left| \sum_{\mathbf{r} \in A} \Psi_{\text{col}}^1(\mathbf{r}) \right|^2 \right\rangle. \quad (4)$$

In a phase with long-range column-phase order we expect $\chi^1 \sim L^{-2}$ for large L , while $\chi^1 \sim O(1)$ in a spin-fluid or staggered phase. In the FS ground state at $V=J$ we find $\chi_{\text{FS}}^1 \sim (\log L)^2$. The results of the numerical calculations of χ^1 for $L=4$ and $L=6$ are shown in Fig. 2. We note that (i) for $V < J$, the strong size dependence of χ^1 and its almost linear dependence upon V/J are consistent with presence of long-range column phase order, and (ii) for $V > J$, the large value of χ^1 for the 4×4 lattice is due to the impossibility of inserting two domains of the staggered phase in this lattice size; the (0,0) winding-number sector is thus strongly frustrated.

A further probe of the nature of the ground state is offered by the quantity g :

$$g = \frac{\left\langle \left| \sum_{\mathbf{r} \in A} \Psi_{\text{col}}^1(\mathbf{r}) \right|^4 \right\rangle}{\left\langle \left| \sum_{\mathbf{r} \in A} \Psi_{\text{col}}^1(\mathbf{r}) \right|^2 \right\rangle^2}, \quad (5)$$

the ratio of the fourth moment of the order parameter to the square of its second moment. In a phase with long-range column-phase order we expect $g \rightarrow 1$ as $L \rightarrow \infty$. However, in a spin-fluid or staggered phase, the fluctuations in Ψ_{col}^1 can be expected to be Gaussian, in which case




FIG. 2. The correlation function χ^1 [Eq. (4)] as a function of V/J .




FIG. 3. The quantity g [Eq. (5)] as a function of V/J .

$g \rightarrow 2$ and $L \rightarrow \infty$. The values of g are shown in Fig. 3. We note (i) the rapid increase in the value of g for $L=6$ near $V/J \approx 1$ as indicating a crossover from a column phase to the staggered phase; (ii) the oscillations in the value of g for $L=6$ and $V > J$ can be traced to (a) changes in the structure of the domain walls as a function of V/J , and (b) the small value of χ^1 whose square appears in the denominator for the expression for g ; (iii) the value of g for the FS state can be calculated to be $g_{\text{FS}}=2$.

The spatial structure of the state can be investigated by the correlation function

$$G(\mathbf{r}) = \langle \Psi_{\text{col}}^{1*}(\mathbf{r}_1) \Psi_{\text{col}}^1(\mathbf{r}_1 + \mathbf{r}) \rangle. \quad (6)$$

For $\mathbf{r}, \mathbf{r}_1 \in A$, the function $G(\mathbf{r})$ can be shown to be real. The results for $G(\mathbf{r})$ are shown in Fig. 4 for $L=6$ and a range of values of V/J . Also shown are the values of $G_{\text{FS}}(\mathbf{r})$, the correlation functions in the FS state. The differences in the values of $G_{\text{FS}}(\mathbf{r})$ and $G(\mathbf{r})$ for $V=J$ in (0,0) winding-number sector (shown in Fig. 5 by the inverted triangles) are due to finite-size corrections. Thus the nonmonotonicity of $G(\mathbf{r})$ at the two largest values of $|\mathbf{r}|$ is a finite-size correction. For $V=0$ and $V=0.5J$, the




FIG. 4. The correlation function $G(\mathbf{r})$ [Eq. (6)] as a function of the Euclidean distance $|\mathbf{r}|$ for various values of V/J on a 6×6 lattice. Also shown are the correlation functions in the infinite lattice FS state.




FIG. 5. The correlation function χ^2 [Eq. (7)] as a function of V/J .

function $G(\mathbf{r})$ clearly appears to asymptote at a nonzero value as $|\mathbf{r}| \rightarrow \infty$, implying long-range column-phase order. The order parameter $\langle \Psi_{\text{col}}^1 \rangle$ satisfies $\lim_{|\mathbf{r}| \rightarrow \infty} G(\mathbf{r}) = |\langle \Psi_{\text{col}}^1 \rangle|^2$: we thus obtain estimates of $|\langle \Psi_{\text{col}}^1 \rangle|$ of 0.53 and 0.40 for $V=0$ and $V=0.5J$, respectively. (Ψ_{col}^1 has been chosen such that a completely ordered rigid column state would have an order-parameter expectation value of 1.)

Finally, we measure x - y symmetry breaking by examining the order parameter Ψ^2 (as $\Psi_{\text{col}}^2 = \Psi_{\text{st}}^2$, we will drop the subscript). We numerically evaluated the correlation function

$$\chi^2 = \frac{1}{L^2} \left\langle \left| \sum_{\mathbf{r} \in A} \Psi^2(\mathbf{r}) \right|^2 \right\rangle. \quad (7)$$

We now expect $\chi^2 \sim L^2$ in both the column and the staggered phases. An intermediate spin-fluid phase would have $\chi^2 \sim O(1)$. The values of χ^2 for $L=4$ and $L=6$ are shown in Fig. 5. The sharp minimum in χ^2 for $L=6$ at $V=J$ is consistent with the restoration of Z_2 symmetry at precisely $V=J$ and nowhere else. The sharp dropoff in the value of χ^2 at $V=1.6J$ for $L=6$ is due to changes in the structure of the domain walls between the different staggered-phase domains; we obtain staggered phases rotated by 90° with respect to each other, which drastically reduces the value of $\sum_{\mathbf{r} \in A} \Psi^2$.

CONCLUSIONS

We have found convincing evidence for the crystallization of the dimers into a state with the symmetry of Fig. 1(a) for $V < 0.5J$. However, the simplest scenario consistent with all of the data is the persistence of the column phase right up to $V=J$. For $V > J$, the model is known to be in the staggered ground state; the point $V=J$ will then be a special singular point displaying power-law dimer correlations. It is notable that none of the data show any explicit signal of the existence of spin-fluid order over a finite range of V . We cannot, however, definitively rule out the existence of the spin-fluid phase over a small, but finite, range of V . Monte Carlo simulations on an 8×8 lattice will therefore be of considerable interest.

ACKNOWLEDGMENTS

I benefited from a very useful discussion with D. Huse on techniques for analyzing the data. I would like to thank D. S. Fisher, G. Grinstein, S. Kivelson, N. Read, and D. Rokhsar for useful conversations. This research was supported in part by the National Science Foundation (NSF) under Grant No. DMR-8857228 and by the Alfred P. Sloan Foundation. It was also supported by NSF Grant No. PHY82-17853, supplemented by funds from NASA.

¹D. Rokhsar and S. Kivelson, *Phys. Rev. Lett.* **61**, 2376 (1988).

²P. W. Anderson, *Science* **235**, 1196 (1987).

³J. G. Bednorz and K. A. Müller, *Z. Phys. B* **64**, 188 (1986); M.

K. Wu *et al.*, *Phys. Rev. Lett.* **58**, 908 (1987).

⁴N. Read and S. Sachdev, *Nucl. Phys. B* **316**, 609 (1989).

⁵N. Read and S. Sachdev, *Phys. Rev. Lett.* **62**, 1694 (1989).

⁶M. E. Fisher and J. Stephenson, *Phys. Rev.* **132**, 1411 (1963).

⁷These and other correlation functions in the FS state are most easily calculated by using the Grassman functional integral method of S. Samuel, *J. Math. Phys.* **21**, 2806 (1980).

⁸E. Dagotto and A. Moreo, *Phys. Rev. D* **31**, 865 (1985); E. R. Gagliano, E. Dagotto, A. Moreo, and F. C. Alcaraz, *Phys. Rev. B* **34**, 1677 (1986).