

Effective lattice models for two-dimensional quantum antiferromagnets

Subir Sachdev and R. Jalabert

*Center for Theoretical Physics, P.O. Box 6666
Yale University, New Haven, CT 06511*

We introduce a 2+1 dimensional lattice model, S_0 , of N complex scalars coupled to a compact $U(1)$ gauge field as a description of quantum fluctuations in $SU(N)$ antiferromagnets. Duality maps are used to obtain a single effective action for the Néel and spin-Peierls order parameters. We examine the phases of S_0 as a function of N : the $N \rightarrow \infty$ limit can be deduced from previous work. At $N = 1$, S_0 describes monopoles and their Berry phases, spin-Peierls order, but not the Néel field: Monte-Carlo simulations show a second-order transition from a spin-Peierls phase to a Higgs phase which is the remnant of the Néel phase.

PACS Nos. 75.10.Jm, 75.50.Ee, 74.65.+n

April, 1990

The destruction of Néel order in $d = 2$ antiferromagnets by zero temperature quantum fluctuations is of great current interest [1]. An intriguing possibility that has emerged from large- N [2, 3] expansions of unfrustrated $SU(N)$ antiferromagnets, and from numerical [4] and series [5] work on weakly frustrated antiferromagnets, is that for certain values of the spin S (all except $S = 2, 4, 6 \dots$ for the square lattice) the disordered phase has a broken lattice symmetry and long-range spin-Peierls order. However questions on the nature of the transition between the Néel and spin-Peierls phase and the possibility of a coexistence phase have not been settled. As a first step towards resolving these issues, we introduce here a lattice effective action S_0 which describes spin fluctuations in $d = 2$ $SU(N)$ quantum antiferromagnets in terms of N complex scalars and a compact $U(1)$ gauge field. S_0 will be shown to have the following properties: *(i)* Using duality we obtain from S_0 an action S_f expressed solely in terms of the physical Néel and spin-Peierls order parameters: S_f is the correct generalization of the popular non-linear sigma ($NL\sigma$) model [6] which focuses exclusively on the Néel order. *(ii)* In the limit $N \rightarrow \infty$, previous results [3] can be used to determine the phases of S_0 ; the results are summarized in Fig 1a. *(iii)* Further duality transforms are used to map S_f onto an effective action, S_d , for monopole-like instantons. *(iv)* At $N = 1$ the Néel order parameter is not present but the form of the monopole interactions, their Berry phases, and the spin-Peierls order parameter survive; this limit is therefore a tractable testing ground for studying the latter effects. Results of Monte-Carlo simulations on S_f for $N = 1$ are summarized in Fig 1b. Finally, we speculate on the implication of these results for the physically relevant case of $N = 2$.

We begin by introducing S_0 as an encapsulation of previous studies [3, 6, 7, 8] of $d = 2$ $SU(N)$ antiferromagnets: we have discretized the imaginary time (τ) direction and the fields

therefore lie on the sites, links and faces of a cubic lattice. We have $S_0 = S_Z + S_A + S_B$ with

$$S_Z = -\frac{N}{2g} \sum_i Z_{i\alpha}^* e^{iA_{i\mu}} Z_{i+\hat{\mu},\alpha} + h.c. ; \quad S_A = \frac{N}{2e^2} \sum_i (\epsilon_{\mu\nu\lambda} \Delta_\nu A_\lambda - 2\pi q_\mu)^2 ; \quad S_B = i \frac{\pi}{2} \sum_{\bar{i}} \zeta_{\bar{i}} \Delta_\mu q_{\bar{i}\mu} \quad (1)$$

The sums over i (\bar{i}) extend over the vertices of the direct (dual) cubic lattice with the i (\bar{i}) dependence of the fields often not explicitly displayed; μ, ν, λ extend over the directions x, y, τ and Δ_μ is the lattice derivative. All fields on the sites or links of the direct (dual) lattice have upper (lower) case letters. S_Z is a lattice version of the CP^{N-1} action [9] with Z_α a N -component complex scalar of fixed unit length ($\sum_\alpha |Z_\alpha|^2 = 1$) and $-\pi < A_\mu < \pi$ a compact $U(1)$ gauge field. In the continuum, the CP^{N-1} model is equivalent to a $NL\sigma$ model which has been previously used [6] as a long wavelength description of $SU(N)$ antiferromagnets. For $N = 2$, Z_α is related to the Néel order parameter \mathbf{n} by $\mathbf{n} = Z_\alpha^* \vec{\sigma}_{\alpha\beta} Z_\beta$ where $\vec{\sigma}$ are the Pauli matrices. S_A is the kinetic energy of A_μ in a periodic gaussian form: q_μ is integer valued field on the links of the dual lattice. Although no such term is present in the bare CP^{N-1} model, it will always be dynamically generated *e.g.* by integrating out the Z_α bosons between momentum scales Λ and $\Lambda/2$ where $\Lambda \sim 1/a$ is the upper cutoff. S_B is the crucial Berry phase [7, 3] for antiferromagnets with spins in the fundamental representation of $SU(N)$ on one sublattice and in the conjugate representation on the other (for $N = 2$ both correspond to spin $S = 1/2$); we will consider this value of the spin exclusively. The *fixed* field $\zeta_{\bar{i}}$ is τ independent and has the values 0, 1, 2, 3 on dual lattice sites with even-even, even-odd, odd-odd, and odd-even spatial co-ordinates (Fig 2). The quantity $\Delta_\mu q_\mu$ is the divergence of the $U(1)$ flux and equals the number of Dirac monopoles. For *small* e^2 and $A_\mu = A_\mu^D$, the vector potential of a Dirac monopole with total flux 2π , the action will be minimized with $q_\mu = 1$ along the Dirac string; $\Delta_\mu q_\mu$ is thus the number operator for the monopoles. The monopoles are closely tied to the ‘hedgehogs’ in the Néel field (see Ref [3] and below) which

were first shown by Haldane [7] to carry Berry phases. For $e^2 \rightarrow \infty$ the fluctuations of q_μ and A_μ are less strongly coupled; in fact at $e^2 = \infty$ the sum over q_μ is trivial and we obtain $\text{Tr}e^{-S_0} = 0$! The large e^2 limit must therefore be taken with some care and we shall show below that the proper Berry phase continues to be associated with each monopole. We will consider properties of S_0 as a function of e^2 and g for all $N \geq 1$; for $N = 1$ S_0 describes a Higgs scalar coupled to a compact $U(1)$ gauge field whose monopoles carry Berry phases. The analyses for all $N \geq 2$ are very similar; we will therefore often present explicit results only for $N = 1, 2$.

Standard duality methods [10] can be used to exchange the summation over q_μ for summation over an integer valued field a_μ lying on the links of the dual lattice. This transforms S_0 into

$$S_1 = \frac{e^2}{2N} \sum_{\bar{i}} \left(a_\mu - \frac{1}{4} \Delta_\mu \zeta \right)^2 + i \sum_i A_\mu \epsilon_{\mu\nu\lambda} \Delta_\nu a_\lambda - \frac{N}{2g} \sum_i Z_{i\alpha}^* e^{iA_{i\mu}} Z_{i+\hat{\mu},\alpha} + h.c. \quad (2)$$

Correlations of the integer-valued dual ‘flux’ [11] $\epsilon_{\mu\nu\lambda} \Delta_\nu a_\lambda$ under S_1 are identical to those of the Z_α boson current under S_0 . Conversely, the $U(1)$ fluxes $(\epsilon_{\mu\nu\lambda} \Delta_\nu A_\lambda - 2\pi q_\mu)$ are dual to $e^2(a_\mu - \Delta_\mu \zeta/4)$; in particular, the spin-Peierls order parameters, Ψ_x, Ψ_y , were shown in Ref. [3] to be proportional to the electric field. We therefore identify [12]

$$\Psi_p = \frac{e^2}{N_s} \epsilon_{pq} \sum_{\bar{i}} (-1)^{\bar{i}_x + \bar{i}_y} \left(a_{\bar{i}q} - \frac{1}{4} \Delta_q \zeta_{\bar{i}} \right) \quad (3)$$

where N_s is the number of sites, $p, q = x, y$, and the four possible signs of the pair Ψ_x, Ψ_y identify the four spin-Peierls states (Fig 2).

The large e^2 limit is straightforward with S_1 . The a_μ fields are frozen at $a_\mu^0 = \aleph(\Delta_\mu \zeta/4)$ where $n = \aleph(x)$ is the integer nearest to x (because $a_\mu^0 \neq \Delta_\mu \zeta/4$, $\text{Tr}e^{-S_1} \propto \exp(-ce^2) \rightarrow 0$ as $e^2 \rightarrow \infty$; this prefactor will however cancel out of all correlation functions). Using the result $\epsilon_{\mu\nu\lambda} \Delta_\nu a_\lambda^0 = \delta_{\mu\tau} (-1)^{i_x + i_y}$ the imaginary term in S_1 becomes $i \sum_i (-1)^{i_x + i_y} A_\tau$. For $A_\mu = A_\mu^D$

the summation over i is exactly that evaluated by Haldane [7] and thus yields the same monopole Berry phase.

It is now useful to introduce the parametrization $\sum_{\alpha} Z_{i\alpha}^* Z_{i+\hat{\mu},\alpha} \equiv Q_{i\mu} \exp(i\mathcal{A}_{i\mu})$ with $Q_{i\mu}$ real, positive and $-\pi < \mathcal{A} < \pi$, We note that for $N = 1$, $Q_{i\mu} = 1$ while for $N = 2$, $Q_{i\mu} = ((1 + \mathbf{n}_i \cdot \mathbf{n}_{i+\hat{\mu}})/2)^{1/2} \approx 1 - a^2(\partial_{\mu}\mathbf{n}_i)^2/8$ where a is the lattice spacing. The functional integral over A_{μ} in S_1 can be performed and yields a product of modified Bessel functions of $Q_{i\mu}/g$. As is conventional [10] the Bessel functions are replaced by their large argument expansion yielding finally (for $N = 2$) the partition function $Z = \int \mathcal{D}\mathbf{n} \sum_{a_{\mu}} e^{-S_f}$ with $S_f = S_a + S_{\mathbf{n}} + S_c$ and

$$\begin{aligned} S_a &= \frac{e^2}{2N} \sum_{\bar{i}} \left(a_{\mu} - \frac{1}{4} \Delta_{\mu} \zeta \right)^2 + \frac{g}{2N} \sum_{\bar{i}} (\epsilon_{\mu\nu\lambda} \Delta_{\nu} a_{\lambda})^2 \\ S_{\mathbf{n}} &= \frac{1}{4ga} \int d^3r (\partial_{\mu} \mathbf{n})^2 + \dots \quad ; \quad S_c = -2\pi i \sum_{\bar{i}} a_{\mu} k_{\mu}(\mathbf{n}) \end{aligned} \quad (4)$$

where the ellipses denote *real* higher-order terms that are generated by the expansion of the Bessel function. The variable $k_{\mu}(\mathbf{n})$ is the topological current of \mathbf{n} [9]

$$k_{\mu}(\mathbf{n}) \equiv \frac{1}{2\pi} \epsilon_{\mu\nu\lambda} \Delta_{\nu} \mathcal{A}_{\lambda} = \frac{1}{4\pi} \Omega(\mathbf{n}_i, \mathbf{n}_{i+\hat{\mu}}, \mathbf{n}_{i+\hat{\mu}+\hat{\nu}}, \mathbf{n}_{i+\hat{\nu}}) \approx \frac{a^2}{4\pi} \epsilon_{\mu\nu\lambda} \mathbf{n} \cdot (\partial_{\nu} \mathbf{n} \times \partial_{\lambda} \mathbf{n}) \quad (5)$$

where $\Omega(\mathbf{n}_1, \mathbf{n}_2, \mathbf{n}_3, \mathbf{n}_4)$ is the area of the spherical cap spanned by $\mathbf{n}_1 \dots \mathbf{n}_4$; the second equality is only true upto an integer which leaves Z unchanged. S_f is the advertised action controlling the fluctuations of the spin-Peierls and Néel order. For $N = 2$ the mapping between S_0 and S_f is similar in spirit to analyses in Ref [13] for the quantum Hall effect. For $N = 1$, there is no Néel order parameter; thus $S_c = S_{\mathbf{n}} = 0$ and $S_f = S_a$.

We now examine the properties of S_0 and S_f in different parameter regimes:

1. $e^2 \rightarrow \infty$: The a_{μ} fields are frozen at $a_{\mu}^0 = \aleph(\Delta_{\mu}\zeta/4)$; S_c can then be shown to equal $i \sum_i (-1)^{i_x+i_y} \mathcal{A}_{\tau}$. This is identical to the individual spin Berry phase found in Ref [7] as, in an appropriate gauge, $\mathcal{A}_{\tau}(i)$ is half the solid angle between $\mathbf{n}(i)$, $\mathbf{n}(i+\hat{\tau})$ and $\hat{\mathbf{z}}$. *Remarkably,*

the monopole Berry phase has uniquely fixed the individual spin Berry phases; the resulting action for \mathbf{n} is thus identical to that of Ref [7]. For $N = 1$, the quenching of a_μ rules out the possibility of a phase transition at $e^2 = \infty$.

2. $e^2 = 0$: The A_μ field is pure gauge and we choose $A_\mu = q_\mu = 0$. The action S_0 is simply that of a *classical* three dimensional $U(N)$ spin model which has a second-order phase transition $g = g_c$ with exponent [14] $\nu = 1/2 + \epsilon(N + 1)/(4(N + 4)) + \dots$ where $\epsilon = 4 - D = 1$. The action $S_f(e^2 = 0)$ is an unconventional representation of this well-known model. Following Peskin [15], it can be shown that [16] the two phases of the $U(N)$ model can be characterized by different asymptotic limits of correlation functions of a_μ alone: the small g phase has a proliferation of macroscopic loops of the Z_α boson current $\epsilon_{\mu\nu\lambda}\Delta_\nu a_\lambda$, which are absent in the large g phase.

3. Small e^2 : The concentration of monopoles is exponentially small ($\sim \exp(-cN/e^2)$); we now show that for *all* values of N oppositely charged monopoles have a bare linear attraction $\sim NR/g$ for small g and a Coulombic $N/(e^2R)$ attraction at large g . The monopole effective interaction is obtained by promoting the sum over a_μ into an integral by the Poisson method; this produces an integer-valued field $v_{\bar{i}\mu}$ residing on the links of the dual lattice which is the vorticity in the Z_α boson current. Integrating over a_μ we get $Z = \int \mathcal{D}\mathbf{n} \sum_{v_\mu} \exp(-S_d)$ with

$$S_d = 2\pi^2 \sum_{\bar{i}, \bar{j}} \left((v_{\bar{i}\mu} + k_{\bar{i}\mu})G(\bar{i} - \bar{j})(v_{\bar{j}\mu} + k_{\bar{j}\mu}) + \frac{g}{e^2} m_{\bar{i}}G(\bar{i} - \bar{j})m_{\bar{j}} \right) + i\frac{\pi}{2} \sum_{\bar{i}} m_{\bar{i}}\zeta_{\bar{i}} + S_{\mathbf{n}} \quad (6)$$

where $G(r)$ is the Green's function of the lattice Helmholtz equation $(-g\Delta^2 + e^2)G(r) = N\delta(r)$. We have used $\Delta_\mu k_\mu = 0$ and introduced the monopole density $m = \Delta_\mu v_\mu$: the monopoles thus terminate lines of vorticity and carry the Berry phase $\pi\zeta/2$. We now examine the v_μ, k_μ fluctuations for a fixed monopole density for large and small g (*i*) *Small g* : Now $G(0) \sim N/g$ is large and for $N = 2$, S_d forces the minimization of $v_\mu + k_\mu$ as in Fig 3; each monopole is thus bound a hedgehog in the \mathbf{n} field. The action for Fig 3 arises predominantly

from $S_{\mathbf{n}}$ and yields [17] to yield a attractive potential $\sim NR/g$ where R is the separation of the monopoles. For $N = 1$, $k_{\mu} = 0$; the action for each monopole-anti-monopole pair is given by that of the v_{μ} string connecting them $\sim G(0)R \sim R/g$. (ii) *Large g* The v_{μ} and k_{μ} fluctuations are less strongly coupled. We integrate out the \mathbf{n} fluctuations in S_f (Eqn 4) by a ‘high-temperature’ expansion and sum over v_{μ} (subject to the constraint $\Delta_{\mu}v_{\mu} = m$) in a Debye-Huckel approximation [18]. This can be shown to yield [16] an effective Coulombic interaction $\sim Nm_{\bar{i}}m_{\bar{j}}/(e^2|r_{\bar{i}} - r_{\bar{j}}|)$ between monopole charges [3] for all N .

4. Large g: The \mathbf{n} fluctuations can be integrated out in a ‘high temperature’ expansion and, apart from an innocuous renormalization of parameters, the physics is independent of N . The results can therefore be deduced from the $N = 1$ case.

We now turn to a detailed study of $N = 1$ for which $S_f = S_a$. For $g = \infty$, a_{μ} is restricted to $a_{\mu} = \Delta_{\mu}\chi$ where χ is an integer-valued scalar field; S_a is now equivalent [3] to the discrete Gaussian model of Ref. [19] obtained via a duality transform on a quantum-dimer representation of the disordered phase of quantum antiferromagnets. Numerical simulations on this model have been carried out [20] and are consistent with our results below. For $g \neq \infty$ it is useful to examine the global minima of S_a . We find two regimes: (i) $g < e^2/4$: The lowest action states have $a_{\mu}^0 = \aleph(\Delta_{\mu}\zeta/4)$ which implies $\Psi_p^0 = 0$. (ii) $g > e^2/4$: Now $a_{\mu}^0 = \Delta_{\mu}\chi^0$. There is however a large degeneracy in the choice of χ^0 with one state being associated with every dimer close-packing of the square lattice.

We have performed Monte-Carlo simulations on S_a to study the effects of fluctuations. The standard Metropolis algorithm was used with 3×10^5 time steps per site. Long equilibration times restricted us to $e^2 \leq 6$. The results are summarized in Fig 1b. We find two phases: (i) a large g spin-Peierls phase where fluctuation effects lead to crystallization of the dimers in columns (Fig 2); at $g = \infty$ spin-Peierls order is expected to exist for all finite

e^2 [3] (ii) a small g Higgs phase in the $U(1)$ scalar Z_α . A scaling plot of the invariant ratio $r = \langle(\Psi_x^2 + \Psi_y^2)^2\rangle/\langle(\Psi_x^2 + \Psi_y^2)\rangle^2$ (Fig 4) shows clearly that the phase transition at $e^2 = 2.0$ is continuous; we find the exponents $\nu = 0.64 \pm 0.05$ and the specific heat peak [16] is consistent with $\alpha \approx 0$. The Z_4 lattice symmetry breaking pattern of the spin-Peierls states suggests that the phase transition is in the universality class of the $D = 3$, Z_4 clock model, or equivalently, the two-color Ashkin Teller model. The second-order transition in the latter model is XY -like [21] and the exponents $\nu_{XY} \approx 2/3$ and $\alpha_{XY} \approx 0$ are consistent with our results. We recall that the phase transition at $e^2 = 0$ is also XY -like but with the small g phase being the ordered one.

The analysis of S_0 and S_f for the important $N = 2$ case is complicated by the imaginary term whose form suggests a ‘dual’ relationship between the Néel and spin-Peierls ordering. We have just argued that for large g , spin-Peierls order will be present. At small g the \mathbf{n} field will order; a spin-wave analysis of S_c shows that \mathbf{n} fluctuations induce a power-law coupling in the a_μ field: $\sim g^2(\vec{\Delta} \times \vec{a})_{i\mu}(1/r^4)(\delta_{\mu\nu} - 3r_\mu r_\nu/r^2)(\vec{\Delta} \times \vec{a})_{j\mu}$ where $r = r_i - r_j$. This coupling will favor fluctuations in the local value of $\vec{\Delta} \times \vec{a}$; the spin-Peierls states have $\vec{\Delta} \times \vec{a} = 0$ and will therefore be suppressed although we cannot rule out coexistence between the Néel and spin-Peierls phases. Conversely, the existence of an intermediate phase with neither Néel or spin-Peierls ordering is not possible at $e^2 = 0$, and has been argued [3] to be unlikely when the monopole concentration ($\sim \exp(-cN/e^2)$ for S_0) is small. A possible scenario is therefore a single phase boundary coming in at a finite value of g as $e^2 \rightarrow \infty$; unlike $N \rightarrow \infty$ (Fig 1a) however, the phase boundary will not be vertical. Monte-Carlo or other analyses of the actions S_0 or S_f for $N = 2$ to resolve these questions is clearly an important subject for future research.

S.S. would like to thank N. Read for a previous collaboration [2, 3] which led to the

present work. We are grateful to C. Camacho, C. Kane, D.H. Lee, S. Liang, N. Read and R. Shankar for useful discussions. S.S. was supported in part by National Science Foundation Grant No. DMR 8857228 and by the Alfred P. Sloan Foundation. R.J. was supported by an IBM fellowship.

References

- [1] S. Chakravarty, B.I. Halperin, and D.R. Nelson, Phys. Rev. Lett., **60**, 1057 (1988).
- [2] N. Read and S. Sachdev, Nucl. Phys. **B316**, 609 (1989).
- [3] N. Read and S. Sachdev, Phys. Rev. Lett., **62**, 1694 (1989); preprint
- [4] E. Dagotto and A. Moreo, Phys. Rev. Lett. **63** 2148 (1989).
- [5] M.P. Gelfand, R.R.P. Singh, and D.A. Huse, AT&T preprint.
- [6] F.D.M. Haldane, Phys. Rev. Lett. **50**, 1153 (1983); I. Affleck, Nucl. Phys. **B257**, 397 (1985); Nucl. Phys. **B265**, 409 (1985).
- [7] F.D.M. Haldane, Phys. Rev. Lett. **61**, 1029 (1988)
- [8] A. Auerbach and D. Arovas, Phys. Rev. **B38**, 316 (1988).
- [9] P. Di Vecchia, A. Holtkamp, R. Musto, F. Nicodemi and R. Pettorino Nucl. Phys. **B190**, 719 (1981); B. Berg and C. Panagiotakopoulos, *ibid.* **B251**, 353 (1985).
- [10] R. Savit, Rev. Mod. Phys. **52**, 453 (1980); C. Dasgupta and B.I. Halperin, Phys. Rev. Lett. **47**, 1556 (1981).
- [11] The action S_1 is not invariant under integer-valued gauge transformations of a_μ for $e^2 \neq 0$: this can be remedied by introducing an integer valued scalar ϕ and changing

the co-efficient of e^2 to $(a_\mu - \Delta_\mu\phi - \Delta_\mu\zeta/4)^2$. The expression (2) for S_1 corresponds to the gauge choice $\phi = 0$ which has been implicitly made in this paper.

[12] W. Zheng and S. Sachdev, Phys. Rev. **B40**, 5204 (1989).

[13] D.H. Lee and C. Kane, Phys. Rev. Lett., **64**, 1313 (1990).

[14] K.G. Wilson, Phys. Rev. Lett. **28**, 548 (1972).

[15] M.E. Peskin, Ann. Phys. **113**, 122 (1978).

[16] R. Jalabert and S. Sachdev, to be published.

[17] S. Ostlund, Phys. Rev. **B24**, 485 (1981).

[18] D.R. Nelson and J. Toner, Phys. Rev. **B24**, 363 (1981).


[19] E. Fradkin and S. Kivelson, Mod. Phys. Lett. **B4**, 225 (1990).

[20] E. Fradkin, S. Kivelson, and S. Liang, private communication.


[21] R.V. Ditzian, J.R. Banavar, G.S. Grest, and L.P. Kadanoff, Phys. Rev. **B22**, 2542 (1980).

Figure Captions

1. Phase diagrams of the effective action S_0 , and its dual S_f . The $N \rightarrow \infty$ result was deduced from Ref [3], the $N = 1$ result from our Monte-Carlo simulations. The phase transition at $e^2 = 0$ is that of a classical three-dimensional $U(N)$ spin model and is thus second-order. For finite e^2 , the transition is second-order *at* $N = \infty$; its behavior for finite but large N is unknown. For $N = 1$ we observed a transition at the points shown (the line is a guide to the eye and is theoretically expected to go into the point $(e^2 = \infty, g = \infty)$); the scaling analysis in Fig 4 for $e^2 = 2.0$ shows that the transition is second-order. In the limit $g \rightarrow \infty$ the model is insensitive to N and is equivalent to the discrete Gaussian model of Fradkin and Kivelson [19] obtained via a duality transform on the quantum dimer model.
2. Fully ordered spin-Peierls state. The numbers denote values of the *fixed* field ζ and the fluctuating field a_μ lies on the arrows. Each dimer covering corresponds to a value of $a_\mu - (\Delta_\mu \zeta)/4$ of $-3/4$ across dimerized links and of $1/4$ across all other links. The particular dimerization shown therefore has $a_\mu = 0$.
3. A pair of point defects at small g showing the binding between monopoles and hedgehogs.
4. Scaling plot of the invariant ratio $r = \langle(\Psi_x^2 + \Psi_y^2)^2\rangle/\langle(\Psi_x^2 + \Psi_y^2)\rangle^2$ at $N = 1$ and $e^2 = 2.0$ for lattices with L^3 sites. As $g \rightarrow \infty$ (the spin-Peierls phase), $r \rightarrow 1$ while as $g \rightarrow 0$, $r \rightarrow 2$ which corresponds to gaussian fluctuations about $\Psi_p = 0$. The fitting parameters are $\nu = 0.64 \pm 0.05$ and $g_c = 2.39$.



(a) $N \rightarrow \infty$



(b) $N = 1$

Figure 1




Figure 2




Figure 3




Figure 4